

STEADY STATES FOR ONE DIMENSIONAL CONFORMAL METRIC FLOWS

YILONG NI AND MEIJUN ZHU

ABSTRACT. We define two conformal structures on S^1 which give rise to a different view of the affine curvature flow and a new curvature flow, the “ Q -curvature flow”. The steady state of these flows are studied. More specifically, we prove four sharp inequalities, which state the existences of the corresponding extremal metrics.

1. INTRODUCTION

The study of geometric flow equations have attracted great attentions recently from scientific community. In its very short history of existence, there have been some very successful and beautiful results. To name a few, we mention the study of Thurston’s geometrization via Ricci flow and the applications of geometric flows in image processing. The philosophy underneath might be that, while mathematicians like “stabilized” manifolds, human vision system *prefers* “stabilized” images, which could be explained as the limits of certain flow processes.

In this paper we shall start to study certain one dimensional geometric flows, which are directly linked to image processing. Our approach however is inspired and strongly influenced by the study of flow equations for high dimensional manifolds, in particular by the study of Yamabe flow for dimension $n \geq 3$.

Recall that on a n -dimensional compact Riemannian manifold (M^n, g) ($n \geq 3$), the conformal Laplacian is defined as

$$(1.1) \quad L_g = \frac{4(n-1)}{n-2} \Delta_g + R_g,$$

where R_g is the scalar curvature of metric g . It is a conformal covariant, namely for $\varphi > 0$,

$$L_{\varphi^{\frac{4}{n-2}}g} u = \varphi^{-\frac{n+2}{n-2}} L_g(\varphi u).$$

The normalized Yamabe flow, so designed as to find a metric with constant scalar curvature in a given conformal class, is defined as

$$(1.2) \quad \partial_t g = (\bar{R}_g - R_g)g,$$

where $\bar{R}_g = \int_M R_g dV_g / \int_M dV_g$. The steady state of (1.2) is a metric with constant scalar curvature. See Ye [15] and recent work by Schwetlick and Struwe [12].

It is obvious that (1.1) is not a well defined differential operator for $n \leq 2$. We shall mimic the above to define a general α -scalar curvature for any given closed convex curve. Let (S^1, g_s) be the unit circle with the induced metric $g_s = d\theta \otimes d\theta$ from R^2 . For any metric g on S^1 (that is, we reparametrize the circle), we write $g := d\sigma \otimes d\sigma = v^{-4} g_s$ for some positive function v and define a general α -scalar

curvature of g for any positive constant α by

$$R_g^\alpha = v^3(\alpha v_{\theta\theta} + v).$$

Thus $R_{g_s}^\alpha = 1$. If we define the corresponding α -conformal Laplace-Beltrami operator of g by

$$L_g^\alpha = \alpha \Delta_g + R_g^\alpha,$$

where $\Delta_g = D_{\sigma\sigma}$, we are able to show that L_g^α is a conformal covariant, and to establish the formula of R_g^α under a conformal change of metric:

Proposition 1. *For $\varphi > 0$, if $g_2 = \varphi^{-4}g_1$ then $R_{g_2}^\alpha = \varphi^3 L_{g_1}^\alpha \varphi$, and*

$$L_{g_2}^\alpha(\psi) = \varphi^3 L_{g_1}^\alpha(\psi\varphi), \quad \forall \psi \in C^2(\mathbf{S}^1).$$

It turns out, among all positive α , two cases of $\alpha = 1$ and $\alpha = 4$ are of special interest.

We first show that the affine curvature of any given convex curve can be viewed as a 1-scalar curvature. To see this, let $\mathbf{x}(\theta)$ ($\theta \in [0, 2\pi]$) be a closed convex curve in $\mathbf{R}^2$. One can introduce a new parameter of the affine arc-length σ by requiring

$$[\mathbf{x}_\sigma, \mathbf{x}_{\sigma\sigma}] = 1.$$

In fact, if $k = k(\theta)$ is the curvature of the curve, one can define

$$\sigma(\theta) = \int_0^\theta k^{-2/3} d\theta = \int_0^s k^{1/3} ds,$$

where s is the parameter of the arc length. Let $v = k^{1/3}$, we have

$$g_1 := d\sigma \otimes d\sigma = v^{-4} d\theta \otimes d\theta = v^{-4} g_s.$$

Then the affine curvature of $\mathbf{x}(\theta)$ is given by:

$$\kappa = v^3(v_{\theta\theta} + v).$$

which coincides with $R_{g_1}^1$ which we defined above. For simplicity we write $\kappa_g = R_g^1$.

The affine flow, which is successfully used in the image processing (see [11]), is defined as

$$\mathbf{x}_t(\sigma, t) = \mathbf{x}_{\sigma\sigma}.$$

It is shown in [11] that the above flow is equivalent to

$$\partial_t g = -\kappa_g g.$$

Further, it will be shown in our forthcoming paper [10] that the affine flow is equivalent to the following one-dimensional normalized affine flow:

$$(1.3) \quad \partial_t g = (\bar{\kappa}_g - \kappa_g)g,$$

where $\bar{\kappa}_g = \int \kappa_g d\sigma / \int d\sigma$; More over, along the normalized flow $\bar{\kappa}_g$ is always increasing. To establish the global existence, as well as to study the long time behaviour of the flow, it is essential to obtain the upper bound for the total curvature $\int \kappa_g d\sigma$ (note that under the normalized flow, total affine length is fixed). In [11], such an upper bound is obtained due to Blaschke-Santaló inequality, which in turn is proved via the affine isoperimetric inequality as well as the classical Minkowski's mixed volume inequality. In this paper, we shall directly establish such an upper bound via a pure analytic argument. In fact, we will give an analytic proof to the

following general Blaschke-Santaló type inequality, which immediately implies that $\bar{\kappa}_g$ is bounded above along the normalized flow.

Theorem 1. (*General Blaschke-Santaló*) For $u(\theta) \in H^1(S^1)$ and $u > 0$, if u satisfies

$$\int_0^{2\pi} \frac{\cos(\theta + \gamma)}{u^3(\theta)} d\theta = 0$$

for all γ , then

$$\int_0^{2\pi} (u_\theta^2 - u^2) d\theta \int_0^{2\pi} u^{-2}(\theta) d\theta \geq -4\pi^2,$$

and the equality holds if and only if

$$u(\theta) = c\sqrt{\lambda^2 \cos^2(\theta - \alpha) + \lambda^{-2} \sin^2(\theta - \alpha)},$$

for some $\lambda, c > 0$ and $\alpha \in [0, 2\pi)$.

In the consequence paper [10], we will show that the global existence of the affine flow (1.3) follows from this inequality and the maximum principle.

For the case of $\alpha = 4$, the 4-scalar curvature R_g^4 can be viewed as the scalar curvature in an analogous one-dimensional Yamabe flow. Let $g = d\delta \otimes d\delta = u^{-4}g_s = u^{-4}d\theta \otimes d\theta$ for some positive function $u \in C^1(S^1)$. We define the scalar curvature k_g of metric g by

$$k_g := u^3(4u_{\theta\theta} + u),$$

and the corresponding conformal Laplace Beltrami operator by

$$(1.4) \quad A_g := 4\Delta_g + k_g,$$

that is $k_g = R_g^4$ and $A_g = L_g^4$. Thus A_g is a conformal covariant by Proposition 1 and

$$k_{g_2} = \varphi^3 A_{g_1} \varphi,$$

for $g_2 = \varphi^{-4}g_1$. The one-dimensional Yamabe flow is guided by

$$(1.5) \quad \partial_t g = -k_g g.$$

It will be shown [10] that this flow is equivalent to the normalized Yamabe flow:

$$(1.6) \quad \partial_t g = (\bar{k}_g - k_g)g,$$

where $\bar{k}_g = \int_g k_g dS_g / \int dS_g$.

The upper bound of $\bar{k}_g$, which is essential to the proof of the global existence of the flow, is given by the following inequality:

Theorem 2. For $u(\theta) \in H^1(S^1)$ and $u > 0$,

$$\int_0^{2\pi} (u_\theta^2 - \frac{1}{4}u^2) d\theta \int_0^{2\pi} u^{-2}(\theta) d\theta \geq -\pi^2,$$

and the equality holds if and only if

$$u(\theta) = c\sqrt{\lambda^2 \cos^2 \frac{\theta - \alpha}{2} + \lambda^{-2} \sin^2 \frac{\theta - \alpha}{2}}$$

for some $\lambda, c > 0$ and $\alpha \in [0, 2\pi)$.

Theorem 2 has been proved in Ai, Chou and Wei [1] and Hang [6], motivated by different problems. Our proof, which is based on a local type inequality (motivated by [7]), is completely different to theirs. See also Remark 4.

The analytic approach to the proofs of Theorem 1 and 2 enables us to study other conformal curvature flows on S^1 involving higher order derivatives. The higher dimensional analogues include the studies of Q -curvature (see, e.g., Fefferman and Graham [5]) and the extremal hermitian metrics (see, e.g. Chen [3]). For any given g on S^1 we write $g = v^{-4/3}g_s$, where g_s is the standard metric. The *symmetric Q curvature* of (S^1, g) is defined as

$$(1.7) \quad Q_g^A = \frac{1}{9}v^{\frac{5}{3}}(v_{\theta\theta\theta\theta} + 10v_{\theta\theta} + 9v),$$

and the corresponding operator P^A is defined as

$$(1.8) \quad P_g^A(f) = \frac{1}{9}\Delta_g^2 f + \frac{10}{9}\nabla_g(\kappa_g \nabla_g f) + Q_g^A f, \quad f \in C^4(\mathbf{x}),$$

where κ_g is the 1-curvature. Similar to Proposition 1, we will show

Proposition 2. (*Conformal covariance of P_g^A*) If $g_2 = \varphi^{-\frac{4}{3}}g_1$, then $Q_{g_2}^A = \varphi^{\frac{5}{3}}P_{g_1}^A\varphi$ and

$$P_{g_2}^A\psi = \varphi^{\frac{5}{3}}P_{g_1}^A(\psi\varphi), \quad \forall \psi \in C^4(S^1).$$

The *symmetric Q -curvature* flow is introduced as

$$(1.9) \quad \partial_t g = (Q_g^A - \overline{Q}_g^A)g,$$

where $\overline{Q}_g^A = \int Q_g^A dS_g / \int dS_g$, along which $\overline{Q}_g^A$ is always decreasing. In fact, we will show that $\overline{Q}_g^A$ is bounded from below by a positive constant.

Theorem 3. For $u(\theta) \in H^2(S^1)$ and $u > 0$, if u satisfies

$$\int_0^{2\pi} \frac{\cos^3(\theta + \alpha)}{u^{5/3}(\theta)} d\theta = 0$$

for all α , then

$$\int_0^{2\pi} (u_{\theta\theta}^2 - 10u_{\theta}^2 + 9u^2) d\theta \left(\int_0^{2\pi} u^{-2/3}(\theta) d\theta \right)^3 \geq C_0 > 0.$$

More over, if u_0 is an extremal function, then the symmetric Q -curvature of $u_0^{-4/3}g_s$ is a constant.

The above might be needed to obtain the global existence and convergence of the symmetric Q -curvature flow. We expect such flow will have interesting applications in image processing (especially in image recognition), see Remark 6 after the proof.

At this stage, it is nature to introduce the one-dimensional Q curvature of (S^1, g) given by

$$(1.10) \quad Q_g = \frac{1}{9}v^{\frac{5}{3}}(16v_{\theta\theta\theta\theta} + 40v_{\theta\theta} + 9v),$$

and the corresponding operator P_g on (S^1, g) given by

$$(1.11) \quad P_g(f) = \frac{16}{9}\Delta_g^2 f + \frac{40}{9}\nabla_g(k_g \nabla_g f) + Q_g f, \quad f \in C^4(S^1),$$

where k_g is the 4-scalar curvature of (S^1, g) . The Q -curvature flow is then guided by

$$(1.12) \quad \partial_t g = (Q_g - \overline{Q}_g)g,$$

where $\overline{Q}_g = \int Q_g dS_g / \int dS_g$. Simliar to Proposition 2, we will show that

Proposition 3. (Conformal Covariant of P_g) If $g_2 = \varphi^{-\frac{4}{3}}g_1$, then $Q_{g_2} = \varphi^{\frac{5}{3}}P_{g_1}\varphi$ and

$$P_{g_2}\psi = \varphi^{\frac{5}{3}}P_{g_1}(\psi\varphi), \quad \forall \psi \in C^4(\mathbf{x}).$$

Remark 1. We may define general $\alpha - Q$ curvature on (S^1, g) as

$$Q_g^\alpha = v^{\frac{5}{3}}\left(\frac{\alpha^2}{9}v_{\theta\theta\theta\theta} + \frac{10\alpha}{9}v_{\theta\theta} + v\right),$$

and the corresponding operator as

$$P_g^\alpha(f) = \frac{\alpha^2}{9}\Delta_g^2 f + \frac{10\alpha}{9}\nabla_g(R_g^\alpha \nabla_g f) + Q_g^\alpha f,$$

where $g = v^{-4/3}g_s$ and R_g^α is the α -scalar curvature of (S^1, g) . Similar to Proposition 3 we have

Proposition 4. (Conformal Covariant of P_g^α) If $g_2 = \varphi^{-\frac{4}{3}}g_1$, then $Q_{g_2}^\alpha = \varphi^{\frac{5}{3}}P_{g_1}^\alpha\varphi$ and

$$P_{g_2}^\alpha\psi = \varphi^{\frac{5}{3}}P_{g_1}^\alpha(\psi\varphi), \quad \forall \psi \in C^4(\mathbf{x}).$$

The following inequality, which classifies the steady state of the flow, is essential to the proof of the global existence and convergence of the flow.

Theorem 4. For $u(\theta) \in H^2(S^1)$ and $u > 0$,

$$\int_0^{2\pi} \left(u_{\theta\theta}^2 - \frac{5}{2}u_\theta^2 + \frac{9}{16}u^2\right)d\theta \left(\int_0^{2\pi} u^{-2/3}(\theta)d\theta\right)^3 \geq 9\pi^4,$$

and the equality holds if and only if

$$u(\theta) = c \left(\lambda^2 \cos^2 \frac{\theta - \alpha}{2} + \lambda^{-2} \sin^2 \frac{\theta - \alpha}{2} \right)^{3/2},$$

for some $\lambda, c > 0$ and $\alpha \in [0, 2\pi)$.

The main theme of this paper is to prove the above four theorems. Note that extremal functions of these four inequalities give the extremal metrics for corresponding conformal curvatures. Therefore we also have classifications of the limiting metrics of the affine flow, Yamabe flow and the Q -curvature flow. The steady state for symmetric Q -curvature flow is not completely clear yet. See Remark 6.

During our writing of this paper, we learned that the analytic approach to the affine isoperimetric as well as the Blasche-Santaló inequalities has been proposed and tried by another group of researchers, see [4]. For example, the analytic proof of Theorem 1 was given in [4]. However, their proof is incomplete (Lemma 6.2 in [4] is not correct since u_0 can vanish on a whole interval). When the paper was in circulation, we were informed by J. Wei [13] that Proposition A.5 in [1] also yields an analytical proof of Theorem 1. The proofs of Theorem 2 and 4 are also obtained

in a recent preprint [6] by Hang. His work seems to be inspired by early work of Yang and Zhu [14] on three dimensional Paneitz operator. Our proofs are different to those in [14] and [6].

The paper is organized as follows. In section 2, we prove the conformal covariance of operators L_g^α , P_g^A and P_g . In section 3, we establish a local sharp inequality, which implies Theorem 1 using the stereographic projection. In section 4, we give a new proof of the Blaschke-Santaló inequality. The idea is also used in the proof of Theorem 3. We prove Theorem 4 and 3 in section 5.

2. CONFORMAL COVARIANCE

Suppose that $g_1 := d\sigma_1 \otimes d\sigma_1 = v^{-4}g_s := v^{-4}d\theta \otimes d\theta$, where g_s is the standard metric on the unit circle. We will use convention that $\Delta_{g_i} = D_{\sigma_i \sigma_i}$ and $\nabla_{g_i} = D_{\sigma_i}$ for $i = 1$ and 2 . Then for $g_2 = \varphi^{-4}g_1 = (\varphi v)^{-4}g_s$, by definition

$$\begin{aligned} R_{g_2}^\alpha &= (\varphi v)^3 (\alpha \Delta_{g_s}(\varphi v) + \varphi v) \\ &= (\varphi v)^3 (\alpha v \Delta_{g_s} \varphi + 2\alpha \nabla_{g_s} \varphi \nabla_{g_s} v + \alpha \varphi \Delta_{g_s} v + \varphi v). \end{aligned}$$

On the other hand,

$$\begin{aligned} \varphi^3 L_{g_1}^\alpha \varphi &= \varphi^3 (\alpha \Delta_{g_1} \varphi + v^3 (\alpha \Delta_{g_s} v + v) \varphi) \\ &= \varphi^3 (\alpha v^2 \nabla_{g_s} (v^2 \nabla_{g_s} \varphi) + \alpha v^3 \varphi \Delta_{g_s} v + v^4 \varphi) \\ &= (\varphi v)^3 (\alpha v \Delta_{g_s} \varphi + 2\alpha \nabla_{g_s} \varphi \nabla_{g_s} v + \alpha \varphi \Delta_{g_s} v + \varphi v). \end{aligned}$$

Therefore $R_{g_2}^\alpha = \varphi^3 L_{g_1}^\alpha \varphi$. It follows that, for any $\psi \in C^2(S^1)$,

$$\begin{aligned} L_{g_2}^\alpha \psi &= \alpha \Delta_{g_2} \psi + R_{g_2}^\alpha \psi = \alpha \varphi^2 \nabla_{g_1} (\varphi^2 \nabla_{g_1} \psi) + \varphi^3 \psi L_{g_1}^\alpha \varphi \\ &= \alpha \varphi^4 \Delta_{g_1} \psi + 2\alpha \varphi^3 \nabla_{g_1} \varphi \nabla_{g_1} \psi + \alpha \varphi^3 \psi \Delta_{g_1} \varphi + R_{g_1}^\alpha \varphi^4 \psi \\ &= \varphi^3 L_{g_1}^\alpha (\psi \varphi). \end{aligned}$$

This completes the proof of Proposition 1.

To prove Proposition 2, we recall that for $g = v^{-4/3}g_s$, the 1-scalar curvature κ_g has the following form:

$$(2.1) \quad \kappa_g = (v^{\frac{1}{3}})^3 ((v^{\frac{1}{3}})_{\theta\theta} + v^{\frac{1}{3}}) = \frac{1}{3} v^{\frac{1}{3}} v_{\theta\theta} - \frac{2}{9} v^{-\frac{2}{3}} (v_\theta)^2 + v^{\frac{4}{3}}.$$

For any $f \in C^4(S^1)$ and $g_1 = v^{-4/3}g_s$,

$$\begin{aligned} \nabla_{g_1} f &= f_\theta v^{\frac{2}{3}}, \\ \Delta_{g_1} f &= f_{\theta\theta} v^{\frac{4}{3}} + \frac{2}{3} v^{\frac{1}{3}} f_\theta v_\theta, \\ \Delta_{g_1}^2 f &= \frac{10}{9} v^{\frac{2}{3}} (2v_\theta^2 f_{\theta\theta} + v_\theta v_{\theta\theta} f_\theta) \\ &\quad + \frac{1}{3} v^{\frac{5}{3}} (3v f_{\theta\theta\theta\theta} + 12v_\theta f_{\theta\theta\theta} + 8v_{\theta\theta} f_{\theta\theta} + 2v_{\theta\theta\theta} f_\theta), \\ \nabla_{g_1} (\kappa_{g_1} \nabla_{g_1} f) &= \nabla_{g_1} (f_\theta (\frac{1}{3} v v_{\theta\theta} - \frac{2}{9} v_\theta^2 + v^2)) \\ &= v^{\frac{2}{3}} (v^2 f_{\theta\theta} - \frac{2}{9} v_\theta^2 f_{\theta\theta} - \frac{1}{9} v_\theta v_{\theta\theta} f_\theta) \\ &\quad + \frac{1}{3} v^{\frac{5}{3}} (v_{\theta\theta} f_{\theta\theta} + 6v_\theta f_\theta + f_\theta v_{\theta\theta\theta}). \end{aligned}$$

Therefore

$$\begin{aligned}
9\varphi^{\frac{5}{3}}P_{g_1}^A(\varphi) &= \varphi^{\frac{5}{3}}(\Delta_{g_1}^2\varphi + 10\nabla_{g_1}(\kappa_{g_1}\nabla_{g_1}\varphi) + 9Q_{g_1}^A\varphi) \\
&= (v\varphi)^{\frac{5}{3}}(v\varphi_{\theta\theta\theta\theta} + 4v_{\theta}\varphi_{\theta\theta\theta} + 6v_{\theta\theta}\varphi_{\theta\theta} + 4v_{\theta\theta\theta}\varphi_{\theta} \\
&\quad + 20v_{\theta}\varphi_{\theta} + 10v\varphi_{\theta\theta}) + 9\varphi^{\frac{5}{3}}Q_{g_1}^A \\
&= (v\varphi)^{\frac{5}{3}}((v\varphi)_{\theta\theta\theta\theta} + 10(v\varphi)_{\theta\theta} + 9v\varphi) \\
&= 9Q_{g_2}^A.
\end{aligned}$$

For any $\psi > 0$, let $g_3 = \psi^{-4/3}g_2$. The above calculation shows that

$$Q_{g_3}^A = \psi^{\frac{5}{3}}P_{g_2}^A\psi.$$

On the other hand, $g_3 = \psi^{-4/3}g_2 = (\varphi\psi)^{-4/3}g_1$, thus

$$Q_{g_3}^A = (\varphi\psi)^{\frac{5}{3}}P_{g_1}^A(\varphi\psi).$$

Comparing the above two equations, we obtain that

$$(2.2) \quad P_{g_2}^A\psi = \varphi^{\frac{5}{3}}P_{g_1}^A(\varphi\psi)$$

for any $\psi > 0$. Finally, for general ψ , choose C sufficiently large, such that $C + \psi > 0$. Applying (2.2) to $C + \psi$ and using the linearity of P_g^A , we obtain that (2.2) holds for all ψ . This completes the proof of Proposition 2. The proof of Proposition 3 is very similar. We shall skip the details.

3. A LOCAL SHARP INEQUALITY

In this section, we shall follow [7] to establish a local sharp inequality, which will yield the global inequality Theorem 2.

For any three positive constants $a, b, r > 0$, We define

$$D_{a,b}(-r, r) = \{f(y) : f(y) - b \in H_0^1(-r, r), f > 0, \int_{-r}^r f^{-2}(y)dy = a\}$$

Proposition 5. *If $a > 2rb^{-2}$, then*

$$\inf_{w \in D_{a,b}(-r, r)} \int_{-r}^r |\nabla w|^2 dy = 2\tau^{1/2} \left(\frac{r}{\lambda} - \arctan\left(\frac{r}{\lambda}\right) \right),$$

where $\tau > 0$ and $\lambda > 0$ are determined by

$$2\tau^{-1/2} \arctan(r\lambda^{-1}) = a, \quad \frac{\lambda^2 + r^2}{\lambda} \tau^{1/2} = b^2.$$

The infimum is achieved if and only if by

$$w(y) = \tau^{1/4} \sqrt{\frac{\lambda^2 + y^2}{\lambda}}.$$

If $a < 2rb^{-2}$, then

$$\inf_{w \in D_{a,b}(-r, r)} \int_{-r}^r |\nabla w|^2 dy = 2\tau^{1/2} \left(-\frac{r}{\lambda} + \operatorname{arctanh}\left(\frac{r}{\lambda}\right) \right),$$

where $\tau < 0$ and $\lambda > r$ are determined by

$$2|\tau|^{-1/2} \operatorname{arctanh}(r\lambda^{-1}) = a, \quad \frac{\lambda^2 - r^2}{\lambda} |\tau|^{1/2} = b^2.$$

The infimum is achieved if and only if by

$$w(y) = |\tau|^{1/4} \sqrt{\frac{\lambda^2 - y^2}{\lambda}}.$$

If $a = 2rb^{-2}$, then

$$\inf_{w \in D_{a,b}(-r,r)} \int_{-r}^r |\nabla w|^2 dy = 0.$$

The infimum is achieved if and only if by $w = b$.

Proof. It is easy to see that: if $a \geq 2rb^{-2}$, then

$$\inf_{w \in D_{a,b}(-r,r)} \int_{-r}^r |\nabla w|^2 dy = \inf_{w \in D_{a,b}(-r,r), w \geq b} \int_{-r}^r |\nabla w|^2 dy;$$

And if $a \leq 2rb^{-2}$,

$$\inf_{w \in D_{a,b}(-r,r)} \int_{-r}^r |\nabla w|^2 dy = \inf_{w \in D_{a,b}(-r,r), w \leq b} \int_{-r}^r |\nabla w|^2 dy.$$

So, if $\{w_i\} \subset D_{a,b}(-r,r)$ is a minimizing sequence satisfying:

$$\lim_{i \rightarrow \infty} \int_{-r}^r |\nabla w_i|^2 dy = \inf_{w \in D_{a,b}(-r,r)} \int_{-r}^r |\nabla w|^2 dy,$$

we may assume without loss of generality (after rearrangement) that $w_i(-y) = w_i(y)$, and when $a > 2rb^{-2}$, $w'_i(y) \leq 0$ for $y > 0$, $w'_i(y) \geq 0$ for $y < 0$; When $a < 2rb^{-2}$, $w'_i(y) \geq 0$ for $y > 0$, $w'_i(y) \leq 0$ for $y < 0$.

Since $\{w_i\}$ is a minimizing sequence, $\int_{-r}^r |\nabla w_i|^2 dy \leq C$ for all i . It follows from $w_i - b \in H_0^1$ that $\|w_i\|_{H^1} \leq C$. Therefore, up to a subsequence,

$$w_i \rightharpoonup w_0 \text{ weakly in } H^1(-r,r).$$

Using Sobolev embedding theorem we obtain that $\|w_i\|_{C^{0,\frac{1}{2}}(-r,r)} \leq C$, and $w_i \rightarrow w_0$ in $C^{0,\alpha}(-r,r)$ for any $\alpha \in (0, \frac{1}{2})$.

Fatou's Lemma implies

$$\int_{-r}^r w_0^{-2}(y) dy \leq \lim_{i \rightarrow \infty} \int_{-r}^r w_i^{-2}(y) dy = a.$$

It follows from $w_0 \in C^{0,\frac{1}{2}}(-r,r)$ that $w_0(y) > 0$ for any $y \in (-r,r)$. Therefore $w_0 \in D_{a,b}(-r,r)$ is a minimizer. It is easy to see that $w_0(y) = w_0(-y)$, $w_0(r) = w_0(-r) = b$, $w_0 \geq b$ if $a > 2b^{-2}$ and $w_0 \leq b$ if $a < 2b^{-2}$. Moreover w_0 satisfies the Euler-Lagrange equation

$$(3.1) \quad w''(y) = \tau w^{-3}.$$

for some constant τ .

If $a > 2rb^{-2}$, it is easy to see that $\tau > 0$. Let

$$(3.2) \quad v_1(y) = \tau^{\frac{1}{4}} \left(\frac{\lambda^2 + y^2}{\lambda} \right)^{\frac{1}{2}}.$$

We can verify that $v_1(y)$ satisfies (3.1), and $v'_1(0) = 0 = w'_0(0)$. Choosing $\lambda = \tau^{-1/2} w_0^2(0)$, we have $v(0) = w_0(0)$. It follows from the uniqueness of the solution of (3.1) that $w_0(y) = v_1(y)$. The boundary condition yields

$$\tau^{\frac{1}{4}} \left(\frac{\lambda^2 + r^2}{\lambda} \right)^{\frac{1}{2}} = b,$$

and $w_0 \in D_{a,b}(-r, r)$ implies

$$a = \int_{-r}^r w_0^{-2} dy = 2\tau^{-1/2} \arctan(r\lambda^{-1}).$$

Thus the infimum is given by

$$\int_{-r}^r |\nabla w_0|^2 dy = 2\tau^{1/2} \left(\frac{r}{\lambda} - \arctan(r\lambda^{-1}) \right).$$

On the other hand, if $w_0 \in D_{a,b}(-r, r)$ is a minimizer, then $w_0 \geq b$ and satisfies (3.1). From the uniqueness of the solution to the ODE, we know $w_0 = v_0$ with $\lambda = \tau^{-1/2} w_0^2(0)$.

If $a < 2rb^{-2}$, then $\tau < 0$. Let

$$v_2(y) = |\tau|^{\frac{1}{4}} \left(\frac{\lambda^2 - y^2}{\lambda} \right)^{\frac{1}{2}}.$$

Again $v_2(y)$ satisfies (3.1), and $v_2'(0) = 0 = w_0'(0)$. We choose $\lambda = |\tau|^{-1/2} w_0^2(0)$, so that $v_2(0) = w_0(0)$. It follows from the uniqueness of the solution of (3.1) that $w_0(y) = v_2(y)$. The boundary condition gives

$$|\tau|^{\frac{1}{4}} \left(\frac{\lambda^2 - r^2}{\lambda} \right)^{\frac{1}{2}} = b,$$

and $w_0 \in D_{a,b}(-r, r)$ implies

$$a = \int_{-r}^r w_0^{-2} dy = 2|\tau|^{-1/2} \operatorname{arctanh}(r\lambda^{-1}).$$

So the infimum is given by

$$\int_{-r}^r |\nabla w_0|^2 dy = 2|\tau|^{1/2} \left(-\frac{r}{\lambda} + \operatorname{arctanh}(r\lambda^{-1}) \right).$$

Similarly, one can see that such v_2 is the only minimizer.

The case of $a = 2rb^{-2}$ is trivial.

□

4. PROOF OF THEOREM 2

Our proof of Theorem 2 is in the same spirit as that of Theorem C in [7]. For $u(\theta) \in H^1(S^1)(\theta \in [-\pi, \pi])$, $u > 0$, without loss of generality (due to rearrangement), we may assume that $u(\theta) = u(-\theta)$ and $u(\theta)$ is decreasing on $[0, \pi]$.

For $\epsilon \in (0, \pi)$, let $S_\epsilon^1 = \{\theta : \theta \in [-\pi + \epsilon, \pi - \epsilon]\}$,

$$r_\epsilon = \tan \frac{\pi - \epsilon}{2} > 0, \text{ and } a_\epsilon = \int_{-\pi + \epsilon}^{\pi - \epsilon} u^{-2}(\theta) d\theta.$$

It is easy to see that $r_\epsilon \rightarrow +\infty$ as $\epsilon \rightarrow 0$, and $\Phi(S_\epsilon^1) = [-r_\epsilon, r_\epsilon]$, where Φ is the stereographic projection $\Phi : S^1 \rightarrow \mathbf{R}^1$, defined as

$$y = \Phi(\theta) = \tan \frac{\theta}{2} \quad \text{for } \theta \in [-\pi, \pi].$$

Let $w(y) = u(\Phi^{-1}(y))\varphi(y)$, where $\varphi(y) = \sqrt{(1+y^2)/2}$. Then we have (note that $\theta = 2 \arctan y$)

$$\begin{aligned} \int_{-r_\epsilon}^{r_\epsilon} w^{-2} dy &= \int_{-r_\epsilon}^{r_\epsilon} u^{-2} (2 \arctan y) \frac{2}{1+y^2} dy \\ &= \int_{-\pi+\epsilon}^{\pi-\epsilon} u^{-2}(\theta) d\theta = a_\epsilon. \end{aligned}$$

Therefore $w(y) \in D_{a_\epsilon, b_\epsilon}(-r_\epsilon, r_\epsilon)$, where $b_\epsilon = w(r_\epsilon) = u(\pi - \epsilon)\sqrt{(1+r_\epsilon^2)/2}$. Using Proposition 5, we have

$$(4.1) \quad \int_{-r_\epsilon}^{r_\epsilon} (w'(y))^2 dy \geq 2\sqrt{\tau} \left(\frac{r_\epsilon}{\lambda} - \arctan \frac{r_\epsilon}{\lambda} \right),$$

where τ and λ are determined by

$$(4.2) \quad 2\tau^{-1/2} \arctan(r_\epsilon \lambda^{-1}) = a_\epsilon, \quad \frac{\lambda^2 + r_\epsilon^2}{\lambda} \tau^{1/2} = b_\epsilon^2.$$

On the other hand,

$$w'(y) = \varphi \frac{d}{dy} u(2 \arctan y) + u(2 \arctan y) \varphi'(y) = u'(\theta) \frac{1}{\varphi(y)} + u(\theta) \varphi'(y),$$

thus

$$\begin{aligned} \int_{-r_\epsilon}^{r_\epsilon} (w'(y))^2 dy &= \int_{-r_\epsilon}^{r_\epsilon} \left[\frac{u_\theta^2}{\varphi^2(y)} + \frac{(u^2)_\theta \varphi'(y)}{\varphi(y)} + u^2 \cdot (\varphi'(y))^2 \right] dy \\ &= \int_{-\pi+\epsilon}^{\pi-\epsilon} u_\theta^2 d\theta - \int_{-r_\epsilon}^{r_\epsilon} \varphi \varphi'' u^2 dy + \varphi \varphi' u^2 (2 \arctan y) \Big|_{-r_\epsilon}^{r_\epsilon} \\ &= \int_{-\pi+\epsilon}^{\pi-\epsilon} \left(u_\theta^2 - \frac{1}{4} u^2 \right) d\theta + b_\epsilon^2 \frac{2r_\epsilon}{1+r_\epsilon^2}. \end{aligned}$$

Therefore (4.1) and (4.2) imply

$$\begin{aligned} \int_{-\pi+\epsilon}^{\pi-\epsilon} \left(u_\theta^2 - \frac{1}{4} u^2 \right) d\theta \int_{-\pi+\epsilon}^{\pi-\epsilon} u^{-2} d\theta &= a_\epsilon \left(\int_{-r_\epsilon}^{r_\epsilon} (w'(y))^2 dy - b_\epsilon^2 \frac{2r_\epsilon}{1+r_\epsilon^2} \right) \\ &\geq a_\epsilon \left(2\sqrt{\tau} \left(\frac{r_\epsilon}{\lambda} - \arctan \frac{r_\epsilon}{\lambda} \right) - b_\epsilon^2 \frac{2r_\epsilon}{1+r_\epsilon^2} \right) \\ &= 4 \arctan \frac{r_\epsilon}{\lambda} \left(\frac{r_\epsilon}{\lambda} - \arctan \frac{r_\epsilon}{\lambda} - \frac{r_\epsilon}{\lambda} \cdot \frac{\lambda^2 + r_\epsilon^2}{1+r_\epsilon^2} \right) \\ &= 4 \arctan \frac{r_\epsilon}{\lambda} \left(\frac{r_\epsilon}{\lambda} \cdot \frac{1-\lambda^2}{1+r_\epsilon^2} - \arctan \frac{r_\epsilon}{\lambda} \right). \end{aligned}$$

Letting $\beta_\epsilon = \arctan \frac{r_\epsilon}{\lambda} \in (0, \frac{\pi}{2})$, and using (4.2), we have

$$a_\epsilon b_\epsilon^2 = 2 \arctan \frac{r_\epsilon}{\lambda} \cdot \frac{\lambda^2 + r_\epsilon^2}{\lambda} = 4r_\epsilon \frac{\beta_\epsilon}{\sin(2\beta_\epsilon)}.$$

Since $b_\epsilon^2 = u^2(\pi - \epsilon)(1 + r_\epsilon^2)/2 = O(r_\epsilon^2)$, and $\lim_{\epsilon \rightarrow 0} a_\epsilon = \int_{-\pi}^{\pi} u^{-2} d\theta := a$, we have $\lim_{\epsilon \rightarrow 0} \beta_\epsilon = \pi/2$. It then follows from (4.2) that $\lim_{\epsilon \rightarrow 0} \lambda = 2\pi a^{-1} u^{-2}(\pi)$. Hence

$$\lim_{\epsilon \rightarrow 0} 4 \arctan \frac{r_\epsilon}{\lambda} \left(\frac{r_\epsilon}{\lambda} \cdot \frac{1-\lambda^2}{1+r_\epsilon^2} - \arctan \frac{r_\epsilon}{\lambda} \right) = 4 \frac{\pi}{2} (0 - \frac{\pi}{2}) = -\pi^2,$$

and

$$\int_{-\pi}^{\pi} (u_{\theta}^2 - \frac{1}{4}u^2) d\theta \int_{-\pi}^{\pi} u^{-2} d\theta \geq -\pi^2.$$

The equality holds if and only if u satisfies the Euler-Lagrange equation

$$u_{\theta\theta} + \frac{1}{4}u = \tau u^{-3}$$

for some positive constant τ . Thus $w(y) = u(\Phi^{-1}(y))\varphi(y) > 0$ satisfies

$$(4.3) \quad w'' = \tau w^{-3}, w(y) = O(|y|), \text{ as } |y| \rightarrow \infty.$$

The first integral of equation (4.3) is

$$(w')^2 = c_1 - \tau w^{-2}.$$

Since $w(y) = O(|y|)$, as $|y| \rightarrow \infty$, we may assume that $w(y)$ attains its minimum at some point a . It is easy to see that $w(y)$ is increasing when $y > a$, decreasing when $y < a$, and all solutions to (4.3) are given by $w(y) = \sqrt{\frac{\tau + c_1^2(y-a)^2}{c_1}}$, that is

$$u(\theta) = c \sqrt{\lambda^2 \cos^2 \frac{\theta - \alpha}{2} + \lambda^{-2} \sin^2 \frac{\theta - \alpha}{2}},$$

where $c = (4\tau)^{\frac{1}{4}}$ and $\lambda = (\tau)^{\frac{1}{4}}/\sqrt{c_1}$. This completes the proof of Theorem 2.

5. A NEW PROOF OF BLASCHKE-SANTALÓ INEQUALITY

The original proof of Blaschke-Santaló inequality and affine isoperimetric inequality involved Minkowski mixed volume inequality for convex body, see for example [9]. Here we shall give a new analytic proof. The analytic approach enables us to extend such inequality to the one involving higher order derivatives (see our Theorem 3), as well as allows us to remove the geometric constraints (i.e. the convexity of the curve). Such an approach was used also in [4] aiming to generalize the affine isoperimetric inequality. However, as we pointed out early that their proof is incomplete.

For any given $\lambda > 0$ and $\theta \in [0, 2\pi]$, let

$$\begin{aligned} \psi_{\lambda}(\theta) &= \sqrt{\lambda^2 \cos^2 \theta + \lambda^{-2} \sin^2 \theta}, \\ \sigma_{\lambda}(\theta) &= \int_0^{\theta} \psi_{\lambda}^{-2} d\theta = \begin{cases} \arctan(\lambda^{-2} \tan \theta), & \theta \in [0, \pi/2] \\ \arctan(\lambda^{-2} \tan \theta) + \pi, & \theta \in (\pi/2, 3\pi/2] \\ \arctan(\lambda^{-2} \tan \theta) + 2\pi, & \theta \in (3\pi/2, 2\pi) \end{cases} \end{aligned}$$

and define

$$(T_{\lambda}u)(\theta) := u(\sigma_{\lambda}(\theta))\psi_{\lambda}(\theta).$$

From conformal invariant properties, we know that

$$(5.1) \quad \int_0^{2\pi} (u_{\theta}^2 - u^2) d\theta = \int_0^{2\pi} ((T_{\lambda}u)_{\theta}^2 - (T_{\lambda}u)^2) d\theta,$$

and

$$(5.2) \quad \int_0^{2\pi} u^{-2} d\theta = \int_0^{2\pi} (T_{\lambda}u)^{-2} d\theta.$$

Also one can easily check that for all $\alpha \in [0, 2\pi)$

$$(5.3) \quad \int_0^{2\pi} \frac{\cos(\theta + \alpha)}{(T_{\lambda}u)^3} d\theta = \sqrt{\lambda^{-2} \cos^2 \alpha + \lambda^2 \sin^2 \alpha} \cdot \int_0^{2\pi} \frac{\cos(\theta + \tilde{\alpha})}{u^3} d\theta,$$

where $\tilde{\alpha} = \sigma_\lambda^{-1}(\alpha)$.

For any $u \in H^1(S^1)$, $u > 0$, let

$$J(u) := \int_0^{2\pi} (u_\theta^2 - u^2) d\theta \cdot \int_0^{2\pi} u^{-2} d\theta.$$

If we denote

$$H_s^1(S^1) := \{u \in H^1(S^1) : u > 0, \int_0^{2\pi} \cos(\theta + \gamma) u^{-3}(\theta) d\theta = 0, \text{ for all } \gamma \in [0, 2\pi)\},$$

it is obvious that

$$\inf_{H_s^1} J(u) \leq J(1) = -4\pi^2 < 0.$$

We first shall prove that $\inf_{H_s^1} J(u)$ is bounded below and the minimizer of $J(u)$ exists.

Suppose $\{u_i\}$ is a sequence of minimizing functions for $\inf_{H_s^1} J(u)$ with $\max_\theta u_i(\theta) = 1$.

Lemma 1. *For each $f \in C^0(S^1)$, there exists a constant a such that $f(a) = f(a + \pi)$.*

Proof. Let $g(\theta) = f(\theta + \pi) - f(\theta)$. Then $g(\theta) = -g(\theta + \pi)$. Intermediate value theorem immediately yields the result. $\square$

Due to Lemma 1, without loss of generality (after rotating), we may assume that $u_i(0) = u_i(\pi)$ and $u_i(\pi/2) \geq u_i(3\pi/2)$. Let

$$\lambda_i = \sqrt{\frac{u_i(\frac{\pi}{2})}{u_i(0)}}, \text{ and } w_i(\theta) = \frac{T_{\lambda_i} u_i}{\|T_{\lambda_i} u_i\|_{L^\infty}}.$$

It is easy to see from (5.1) to (5.3) that $\{w_i\}$ is also a minimizing sequence and

$$w_i(0) = w_i(\frac{\pi}{2}) = \frac{\sqrt{u_i(0)u_i(\frac{\pi}{2})}}{\|T_{\lambda_i} u_i\|_{L^\infty}}.$$

Furthermore $u_i(0) = u_i(\pi)$ implies $w_i(\pi) = w_i(0) = w_i(\pi/2)$. Since $\{w_i\}$ is a minimizing sequence, we have

$$\int_0^{2\pi} (w_i)_\theta^2 - (w_i)^2 d\theta \leq 0$$

for large i . It follows from $\|w_i\|_{L^\infty} = 1$ that $\|w_i\|_{H^1} \leq C$. Hence $w_i \rightharpoonup w_0$ in H^1 . From Sobolev embedding theorem we have that $w_i \rightarrow w_0$ in $C^{0,\alpha}$ for any $\alpha \in (0, \frac{1}{2})$. If $\min w_0 > 0$, then we know that $\inf_{H_s^1} J(u)$ is bounded below and w_0 is a minimizer. Therefore we only need to consider the case when $\min w_0 = 0$.

We need the following lemma to locate the zero points of w_0 .

Lemma 2. *If $I = (a, b)$ is an open interval satifying $w_0(a) = w_0(b) = 0$ and $w_0(\theta) > 0$ in (a, b) , then $b - a = \pi$.*

Proof. If $b - a > \pi$, let $\delta = b - a - \pi$ and $J = (a + \frac{\delta}{3}, b - \frac{\delta}{3})$. Then $|J| = b - a - \frac{2\delta}{3} > \pi$. Therefore we can choose $\beta > 0$ such that $\cos(\theta + \beta) > 0$ for $\theta \in S^1 \setminus J$, thus

$$\int_{S^1 \setminus J} \frac{\cos(\theta + \beta)}{w_i^3} d\theta \rightarrow \infty \text{ as } i \rightarrow \infty,$$

while $\int_J \frac{\cos(\theta+\beta)}{w_i^3} d\theta$ stays bounded. So, as $i \rightarrow \infty$

$$\int_0^{2\pi} \frac{\cos(\theta+\beta)}{w_i^3} d\theta \rightarrow \infty.$$

This contradicts with

$$\int_0^{2\pi} \frac{\cos(\theta+\beta)}{w_i^3} d\theta = 0.$$

On the other hand, we write the open set $\{\theta : w_0(\theta) > 0\}$ as a union of open intervals

$$\{\theta : w_0(\theta) > 0\} = \cup_i I_i.$$

From the above argument we know that $|I_i| \leq \pi$ for all i . If one of these intervals, I_{i_0} has length strictly less than π , then

$$\int_{I_{i_0}} ((w'_0)^2 - w_0^2) d\theta > 0,$$

and

$$\int_0^{2\pi} ((w'_0)^2 - w_0^2) d\theta \geq \int_{w_0 > 0} ((w'_0)^2 - w_0^2) d\theta = \sum_i \int_{I_i} ((w'_0)^2 - w_0^2) d\theta > 0.$$

It follows that $J(w_i) > 0$ for large i . This contradicts with the fact that

$$\inf_{H_s^1} J(u) \leq J(1) = -4\pi^2.$$

□

Suppose $w_0(\theta_0) = \min w_0 = 0$ for some $\theta_0 \in [0, \pi]$. Since $w_0(0) = w_0(\pi) = w_0(\pi/2) \geq w_0(3\pi/2)$, the above lemma implies that if $w_0(0) = 0$, then $w_0 \equiv 0$. This contradicts with the fact of $\max w_0 = 1$. It follows that $\theta_0 \neq 0$, $w_0(\theta_0) = w_0(\theta_0 + \pi) = 0$ and $w_0(\theta) > 0$ for $\theta \neq \theta_0, \theta_0 + \pi$.

For each i , if $w_i(\theta_i) = \min w_i(\theta)$, then up to a subsequence, we may assume, without loss of generality, that $\theta_i \rightarrow \theta_0 + \pi$. Furthermore, by rotating we may assume that $\theta_0 = 0$, and $\theta_i = \theta_0 + \pi = \pi$, that is, w_i attains its minimum at π , $w_i \rightarrow w_0$ and $w_0 > 0$ for $\theta \in [0, 2\pi] \setminus \{0, \pi\}$.

Let

$$\tau_i = \sqrt{\frac{w_i(\frac{\pi}{2})}{w_i(0)}}, \quad \sigma_i(\theta) = \sigma_{\tau_i}(\theta), \quad \text{and} \quad v_i = \frac{T_{\tau_i} w_i}{\|T_{\tau_i} w_i\|_{L^\infty}}.$$

Again $\{v_i\}$ is a minimizing sequence and $\|v_i\|_{H^1} \leq C$. Therefore, up to a subsequence, $\{v_i\}$ converges weakly in H^1 to some $v_0 \in H^1$ and $v_i \rightarrow v_0$ in $C^{0,\alpha}$ for any $\alpha \in (0, \frac{1}{2})$.

It is easy to see that

$$v_i(0) = v_i(\frac{\pi}{2}), \quad \lim_{i \rightarrow \infty} \frac{v_i(\frac{3\pi}{2})}{v_i(\frac{\pi}{2})} = \frac{w_0(\frac{3\pi}{2})}{w_0(\frac{\pi}{2})} \leq 1, \quad \text{and} \quad v_i(\pi) \leq v_i(0).$$

Hence

$$(5.4) \quad v_0(0) = v_0(\frac{\pi}{2}), \quad \frac{v_0(\frac{\pi}{2})}{v_0(\frac{3\pi}{2})} \geq 1 \quad \text{and} \quad v_0(\pi) \leq v_0(0).$$

Applying Lemma 2 to v_0 , we see that $v_0(0) > 0$, otherwise $v_0(0) = v_0(\frac{\pi}{2}) = v_0(\pi) = v_0(\frac{3\pi}{2}) = 0$, which yields that $v_0 \equiv 0$ by Lemma 2, contradiction to $\max v_0 = 1$. It follows from (5.4) that $v_0(\frac{\pi}{2}) \neq 0$, thus $v_0(\frac{3\pi}{2}) \neq 0$. Applying Lemma 2 again, we

have $v_0(\pi) > 0$. For any $\theta \in [0, 2\pi] \setminus \{\frac{\pi}{2}, \frac{3\pi}{2}\}$, since w_i attains its minimum at π , we obtain

$$\begin{aligned} v_i(\theta) &= \frac{1}{\|T_{\tau_i} w_i\|_{L^\infty}} w_i(\sigma_i(\theta)) \sqrt{\tau_i^2 \cos^2 \theta + \tau_i^{-2} \sin^2 \theta} \\ &\geq \frac{1}{\|T_{\tau_i} w_i\|_{L^\infty}} w_i(\pi) \tau_i |\cos \theta| \\ &= v_i(\pi) |\cos \theta| \rightarrow v_0(\pi) |\cos \theta| > 0 \end{aligned}$$

Hence $v_0 > 0$, thus it is a minimizer.

To complete the proof, we suppose that v is a minimizer and observe that

$$H_s^1(S^1) = \{u \in H^1(S^1) : u > 0, \int_0^{2\pi} \cos \theta \cdot u^{-3} d\theta = \int_0^{2\pi} \sin \theta \cdot u^{-3} d\theta = 0\}.$$

Thus $v(\theta)$ satisfies the Euler-Lagrange equation:

$$(5.5) \quad v_{\theta\theta} + v = \tau v^{-3} + \frac{a \cos \theta}{v^4} + \frac{b \sin \theta}{v^4} = \tau v^{-3} + \frac{c \cos(\theta + \alpha)}{v^4},$$

where τ, a, b are Lagrange multipliers, $c = \sqrt{a^2 + b^2}$ and $\gamma = \gamma(a, b)$. Multiplying both sides of (5.5) by $\cos(\theta + \gamma)$ and integrating over $[0, 2\pi]$, we have $c = 0$. Hence, v satisfies

$$v_{\theta\theta} + v = \tau v^{-3}.$$

Due to Lemma 1, we may assume $v(0) = v(\pi)$. Define w on S^1 by $w(\theta) := v(\frac{\theta}{2})$, $\theta \in [0, 2\pi]$. One can check that $w \in H^1(S^1)$ satisfies

$$w_{\theta\theta} + \frac{1}{4}w = \frac{1}{4}\tau w^{-2}, \quad \theta \in S^1$$

Using the same argument as in the proof of Theorem 1, we obtain that

$$w(\theta) = (\tau)^{1/4} \sqrt{\lambda^2 \cos^2 \frac{\theta - \alpha_1}{2} + \lambda^{-2} \sin^2 \frac{\theta - \alpha_1}{2}}.$$

Therefore,

$$v(\theta) = c \sqrt{\lambda^2 \cos^2(\theta - \alpha) + \lambda^{-2} \sin^2(\theta - \alpha)}.$$

Remark 2. Our argument also yields another proof of the affine isoperimetric inequality. In this regard, we refer [4] to interesting readers for the setting up.

6. HIGHER ORDER DERIVATIVES

We shall prove Theorem 3 and 4 in this section.

We first prove Theorem 4. For any given $\lambda > 0$ and $\alpha \in [0, 2\pi]$, let

$$\Psi_{\lambda, \alpha}(\theta) = \left(\lambda^2 \cos^2 \frac{\theta - \alpha}{2} + \lambda^{-2} \sin^2 \frac{\theta - \alpha}{2} \right)^{3/2},$$

$$\omega_{\lambda, \alpha}(\theta) = \alpha + \int_{\alpha}^{\theta} \Psi_{\lambda, \alpha}^{-\frac{2}{3}} d\theta = \begin{cases} \alpha + 2 \arctan(\lambda^{-2} \tan \frac{\theta - \alpha}{2}), & \text{if } 0 \leq \theta - \alpha \leq \pi \\ \alpha + 2 \arctan(\lambda^{-2} \tan \frac{\theta - \alpha}{2}) + 2\pi, & \text{if } \pi < \theta - \alpha \leq 2\pi \end{cases}$$

and

$$(\mathcal{T}_{\lambda, \alpha} u)(\theta) = u(\omega_{\lambda, \alpha}(\theta)) \Psi_{\lambda, \alpha}(\theta).$$

From conformal invariant properties, we know that

$$\int_0^{2\pi} (u_{\theta\theta}^2 - \frac{5}{2}u_\theta^2 + \frac{9}{16}u^2)d\theta = \int_0^{2\pi} ((\mathcal{T}_{\lambda,\alpha}u)_{\theta\theta}^2 - \frac{5}{2}(\mathcal{T}_{\lambda,\alpha}u)_\theta^2 + \frac{9}{16}(\mathcal{T}_{\lambda,\alpha}u)^2)d\theta$$

and

$$\int_0^{2\pi} u^{-2/3}d\theta = \int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)^{-2/3}d\theta.$$

Lemma 3. *For any $u \in H^2(S^1)$, $u > 0$, there exist $\alpha \in [0, 2\pi)$ and $\lambda \geq 1$, such that*

$$\int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)(\theta) \cos \theta d\theta = \int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)(\theta) \sin \theta d\theta = 0.$$

Proof. For $\alpha \in S^1$, $\lambda \geq 1$, define

$$I_{\lambda,\alpha}(u) = \left(\int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)(\theta) \cos \theta d\theta, \int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)(\theta) \sin \theta d\theta \right) \in \mathbf{R}^2.$$

Suppose $\exists u > 0$, such that $I_{\lambda,\alpha} \neq (0, 0)$ for all $(\lambda, \alpha) \in [1, \infty) \times S^1$. Define $G : [1, \infty) \times S^1 \rightarrow S^1$ by

$$G(\lambda, \alpha) = \frac{I_{\lambda,\alpha}(u)}{\|I_{\lambda,\alpha}(u)\|_{\mathbf{R}^2}},$$

where $\|\mathbf{X}\|_{\mathbf{R}^2}$ is the length of vector $\mathbf{X} \in \mathbf{R}^2$. One can easily check that G is continuous.

When $\lambda = 1$, $\omega_{1,\alpha}(\theta) = \theta$, $\Psi_{1,\alpha}(\theta) = 1$. Therefore

$$\begin{aligned} \int_0^{2\pi} (\mathcal{T}_{1,\alpha}u)(\theta) \cos \theta d\theta &= \int_0^{2\pi} u(\theta) \cos \theta d\theta, \\ \int_0^{2\pi} (\mathcal{T}_{1,\alpha}u)(\theta) \sin \theta d\theta &= \int_0^{2\pi} u(\theta) \sin \theta d\theta. \end{aligned}$$

That is $G(1, \cdot)$ maps S^1 to single point on S^1 .

On the other hand, as $\lambda \rightarrow \infty$, $\omega(\theta) \rightarrow \alpha$ pointwisely for $\theta \in [\alpha, \alpha + \pi)$ and $\omega(\theta) \rightarrow \alpha + 2\pi$ pointwisely for $\theta \in (\alpha + \pi, \alpha + 2\pi]$. Therefore as $\lambda \rightarrow \infty$

$$\begin{aligned} \lambda^{-3} \int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)(\theta) \cos \theta d\theta &= \lambda^{-3} \int_0^{2\pi} u(\omega_{\lambda,\alpha}(\theta)) \Psi_{\lambda,\alpha}(\theta) \cos \theta d\theta \\ &= \int_0^{2\pi} u(\omega_{\lambda,\alpha}(\theta)) \left(\cos^2 \frac{\theta - \alpha}{2} + \frac{1}{\lambda^4} \sin^2 \frac{\theta - \alpha}{2} \right)^{\frac{3}{2}} \cos \theta d\theta \\ &\rightarrow \frac{8}{5} u(\alpha) \cos \alpha \end{aligned}$$

and

$$\lambda^{-3} \int_0^{2\pi} (\mathcal{T}_{\lambda,\alpha}u)(\theta) \sin \theta d\theta \rightarrow \frac{8}{5} u(\alpha) \sin \alpha.$$

Hence

$$\lim_{\lambda \rightarrow \infty} G(\lambda, \alpha) = (\cos \alpha, \sin \alpha).$$

That is $G(\infty, \cdot) = \text{Id}$, the identity map. Since $\pi_1(S^1) \neq 0$, Id is not homotopy to the constant map. Contradiction. $\square$

Remark 3. *There are some similarities between the above lemma and other rearrangement lemma (see, for example, Lemma 2 on Page 85 in [2]).*

For any $u \in H^1(S^1)$, $u > 0$, let

$$F(u) := \int_0^{2\pi} (u_{\theta\theta}^2 - \frac{5}{2}u_\theta^2 + \frac{9}{16}u^2)d\theta \cdot \left(\int_0^{2\pi} u^{-2/3}d\theta \right)^3.$$

If $\{u_i\}$ is a minimizing sequence of F with $\int_0^{2\pi} u_i^{-2/3}d\theta = 1$, due to the above lemma, we may assume, without loss of generality, that u_i satisfies

$$\int_0^{2\pi} u_i \cos \theta d\theta = \int_0^{2\pi} u_i \sin \theta d\theta = 0.$$

Consider the Fourier expansion of u_i :

$$u_i(\theta) = c_{i,0} + \sum_{k=2}^{\infty} (a_{i,k} \cos k\theta + b_{i,k} \sin k\theta).$$

It follows that

(6.1)

$$\begin{aligned} F(u_i) &= \pi \sum_{k=2}^{\infty} k^4 (a_{i,k}^2 + b_{i,k}^2) - \frac{5\pi}{2} \sum_{k=2}^{\infty} k^2 (a_{i,k}^2 + b_{i,k}^2) + \frac{9\pi}{8} c_{i,0}^2 + \frac{9\pi}{16} \sum_{k=2}^{\infty} (a_{i,k}^2 + b_{i,k}^2) \\ &\geq \frac{\pi}{4} \sum_{k=2}^{\infty} k^4 (a_{i,k}^2 + b_{i,k}^2) + \frac{9\pi}{8} c_{i,0}^2 \geq C \int_0^{2\pi} ((u_i)_{\theta\theta})^2 + u_i^2 d\theta \end{aligned}$$

for some positive constant C . Since $\{u_i\}$ is a minimizing sequence with $\int_0^{2\pi} u_i^{-2/3}d\theta = 1$, $F(u_i)$ is bounded. It follows that $\|u_i\|_{H^2} \leq C$. Thus $u_i \rightharpoonup u_0$ weakly in $H^2(S^1)$. Using Sobolev embedding theorem we obtain that $u_0, u_i \in C^{1, \frac{1}{2}}(S^1)$ and $u_i \rightarrow u_0$ in $C^{1, \alpha}(S^1)$ for any $\alpha \in (0, \frac{1}{2})$.

If u_0 vanishes at some point, then $\int_0^{2\pi} u_0^{-2/3}d\theta = \infty$ since $u_0 \in C^{1, \frac{1}{2}}(S^1)$. On the other hand, Fatou's Lemma implies

$$\int_0^{2\pi} u_0^{-2/3}(\theta)d\theta \leq \liminf_{i \rightarrow \infty} \int_0^{2\pi} u_i^{-2/3}(\theta)d\theta = 1.$$

Contradiction. Therefore $u_0 > 0$, thus it is a minimizer.

Suppose u is a minimizer, then it satisfies the Euler-Lagrange equation

$$u_{\theta\theta\theta\theta} + \frac{5}{2}u_{\theta\theta} + \frac{9}{16}u = \tau u^{-\frac{5}{3}},$$

where τ is a positive constant, since $\inf F(u)$ is positive (by (6.1)). Using the Green's function of the operator P_g , we obtain that

$$(6.2) \quad u(\theta) = c \int_0^{2\pi} \tau u^{-\frac{5}{3}}(\varphi) \left| \sin \frac{\theta - \varphi}{2} \right|^3 d\varphi.$$

Define $v : \mathbf{R} \rightarrow \mathbf{R}$ by

$$(6.3) \quad v(y) := u(\Phi^{-1}(y))\varphi^3(y),$$

where $\Phi : S^1 \rightarrow \mathbf{R}^1$ is the stereographic projection:

$$\Phi(\theta) = \tan \frac{\theta}{2}.$$

and

$$\varphi(y) := \left(\frac{1 + y^2}{2} \right)^{1/2}.$$

It follows from (6.2) that $v(y)$ satisfies

$$v(y) = c \int_{-\infty}^{\infty} \tau v^{-\frac{5}{3}} |x - y|^3 dx.$$

Using Theorem 1.5 in Li [8] we obtain that

$$v(y) = c \left(\frac{\lambda^2 + \lambda^{-2}(y-a)^2}{2} \right)^{3/2},$$

for some $c, \lambda > 0$. Therefore

$$u(\theta) = c \left(\lambda^2 \cos^2 \frac{\theta - \alpha}{2} + \lambda^{-2} \sin^2 \frac{\theta - \alpha}{2} \right)^{3/2}.$$

This completes the proof of Theorem 4.

Remark 4. *The same argument yields another proof of Theorem 2.*

Remark 5. *Theorem 4 implies that (S^1, g) has constant Q curvature if and only if it has constant 4-scalar curvature.*

Finally, we shall turn to the proof of Theorem 3. For any $\lambda > 0$, let

$$\Gamma_\lambda(\theta) = (\lambda^2 \cos^2 \theta + \lambda^{-2} \sin^2 \theta)^{3/2}, \quad \sigma_\lambda(\theta) = \int_0^\theta \Gamma_\lambda^{-2/3} d\theta,$$

and

$$(\mathbf{T}_\lambda u)(\theta) := u(\sigma_\lambda(\theta)) \Gamma_\lambda(\theta).$$

By conformal invariant property, we know that

$$\int_0^{2\pi} (u_{\theta\theta}^2 - 10u_\theta^2 + 9u^2) d\theta = \int_0^{2\pi} ((\mathbf{T}_\lambda u)_{\theta\theta}^2 - 10(\mathbf{T}_\lambda u)_\theta^2 + 9(\mathbf{T}_\lambda u)^2) d\theta$$

and

$$\int_0^{2\pi} u^{-2/3} d\theta = \int_0^{2\pi} (\mathbf{T}_\lambda u)^{-2/3} d\theta.$$

Also one can easily check: for any $\alpha \in [0, 2\pi)$,

$$\int_0^{2\pi} \frac{\cos^3(\theta + \alpha)}{(\mathbf{T}_\lambda u)^{5/3}} d\theta = (\lambda^{-2} \cos^2 \alpha + \lambda^2 \sin^2 \alpha)^{3/2} \int_0^{2\pi} \frac{\cos^3(\theta + \tilde{\alpha})}{u^{5/3}} d\theta,$$

where $\tilde{\alpha} = \sigma_\lambda^{-1}(\alpha)$. For any $u \in H^2(S^1)$, $u > 0$, let

$$\mathbf{F}(u) := \int_0^{2\pi} (u_{\theta\theta}^2 - 10u_\theta^2 + 9u^2) d\theta \cdot \left(\int_0^{2\pi} u^{-2/3} d\theta \right)^3,$$

and

$$H_s^2(S^1) := \{u \in H^2(S^1) : \int_0^{2\pi} \frac{\cos^3(\theta + \alpha)}{u^{5/3}} d\theta = 0 \text{ for all } \alpha \in [0, 2\pi)\}.$$

We first prove the existence of a minimizer of $\inf_{H_s^2(S^1)} \mathbf{F}(u)$.

Suppose that $\{u_i\}$ is a minimizing sequence of $\inf_{H_s^2(S^1)} \mathbf{F}(u)$, with $\|u_i\|_{L^\infty} = 1$. The following lemma implies $\|u_i\|_{H^2}$ is bounded, therefore $u_i \rightharpoonup u_0$ in $H^2(S^1)$. Using Sobolev embedding theorem we obtain that $u_i, u_0 \in C^{1, \frac{1}{2}}$, and $u_i \rightarrow u_0$ in $C^{1, \alpha}$, for any $\alpha \in (0, \frac{1}{2})$.

Lemma 4. *Suppose $\{v_i\}$ is a minimizing sequence of $\inf_{H_s^2(S^1)} \mathbf{F}$ with $\|v_i\|_{L^\infty} = 1$. Then $\|v_i\|_{H^2}$ is bounded.*

Proof. We argue by contradiction. Suppose that $\|v_i\|_{H^2}$ is unbounded. Up to a subsequence, we may assume that $\lim_{i \rightarrow \infty} \|v_i\|_{H^2} = \infty$. Let $w_i = v_i / \|v_i\|_{H^2}$. Then $\{w_i\}$ is also a minimizing sequence, $w_i \rightharpoonup 0$ in $H^2(S^1)$ and $\|w_i\|_{H^2} = 1$. It follows that $w_i \rightarrow 0$ in $C^{1,\alpha}$ for any $\alpha \in (0, \frac{1}{2})$, thus $\lim_{i \rightarrow \infty} \int_0^{2\pi} (w_i^2 + (w_i)_\theta^2) d\theta = 0$. On the other hand, we know from $w_i \rightharpoonup 0$ in $H^2(S^2)$ that

$$\lim_{i \rightarrow \infty} \int_0^{2\pi} (w_i)^{-2/3} d\theta = \infty.$$

Note that

$$\lim_{i \rightarrow \infty} \int_0^{2\pi} ((w_i)_{\theta\theta}^2 - 10(w_i)_\theta^2 + 9w_i^2) d\theta \geq 0,$$

and $\{w_i\}$ is a minimizing sequence. We thus have

$$\lim_{i \rightarrow \infty} \int_0^{2\pi} ((w_i)_{\theta\theta}^2 - 10(w_i)_\theta^2 + 9w_i^2) d\theta = 0.$$

Therefore $\lim_{i \rightarrow \infty} \int_0^{2\pi} ((w_i)_{\theta\theta}^2) d\theta = 0$. Hence $\lim_{i \rightarrow \infty} \|w_i\|_{H^2} = 0$. This contradicts with $\|w_i\|_{H^2} = 1$. $\square$

If $u_0 > 0$, we know as in the proof of Theorem 1 that u_0 is a minimizer of $\inf_{H_s^2(S^1)} \mathbf{F}(u)$. The subtle case is that $\min u_0 = 0$.

We need one more lemma.

Lemma 5. *If $I = (a, b)$ is an open interval satisfying $u_0(a) = u_0(b) = 0$ and $u_0(\theta) > 0$ in (a, b) , then $b - a = \pi$.*

Proof. If $b - a > \pi$, let $\delta = b - a - \pi$ and $J = (a + \frac{\delta}{3}, b - \frac{\delta}{3})$. Then $|J| = b - a - \frac{2\delta}{3} > \pi$. Therefore we may choose $\beta > 0$ such that $\cos(\theta + \beta) > 0$ for $\theta \in S^1 \setminus J$. Since $u_0 \in C^{1, \frac{1}{2}}$,

$$\int_{S^1 \setminus J} \frac{\cos^3(\theta + \beta)}{u_i^{5/3}} d\theta \rightarrow \infty, \text{ as } i \rightarrow \infty,$$

while $\int_J \cos^3(\theta + \beta) / u_i^{5/3} d\theta$ is bounded. Therefore, as $i \rightarrow \infty$

$$\int_0^{2\pi} \frac{\cos^3(\theta + \beta)}{u_i^{5/3}} d\theta \rightarrow \infty.$$

Contradiction with the fact that $u_i \in H_s^2(S^1)$.

On the other hand, write the open set $\{\theta : u_0(\theta) > 0\}$ as a union of open intervals:

$$\{\theta : u_0(\theta) > 0\} = \cup_i I_i.$$

From the above argument we know that $|I_i| \leq \pi$ for all i . We claim that for interval I_k with $|I_k| \leq \pi$

$$\int_{I_k} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta \geq 0,$$

and for I_l with $|I_l| < \pi$,

$$\int_{I_l} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta > 0.$$

Therefore, if for some i_0 , $|I_{i_0}| < \pi$, we have

$$\begin{aligned} \int_0^{2\pi} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta &\geq \int_{u_0 > 0} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta \\ &= \sum_i \int_{I_i} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta \\ &\geq \int_{I_0} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta = C > 0. \end{aligned}$$

Since $u_0 \in C^{1,1/2}$, thus $\int_0^{2\pi} u_i^{-2/3} d\theta \rightarrow \infty$. It follows that

$$\lim_{i \rightarrow \infty} \mathbf{F}(u_i) = \infty.$$

Contradiction.

We return to the above claim. Suppose $|I_k| = \pi$. Without loss of generality we may assume that $I_k = (0, \pi)$, i.e. $u_0(0) = 0$, $u_0(\pi) = 0$ and $u_0(\theta) > 0$ when $\theta \in (0, \pi)$. Since $u \geq 0$ and $u \in C^{1, \frac{1}{2}}$, we obtain that $u'(0) = u'(\pi) = 0$. Therefore if $v(\theta) = u(\frac{\theta}{2})$ for $\theta \in [0, 2\pi]$, then $v \in H^2(S^1)$. It follows from Theorem 4 that

$$\int_{I_k} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta = 16 \int_0^{2\pi} (v_{\theta\theta}^2 - \frac{5}{2}v_\theta^2 + \frac{9}{16}v^2) d\theta \geq 0.$$

For $|I_l| < \pi$, we assume that $I_l = (-a, a)$ with $a < \frac{\pi}{2}$. Let $v(\theta) = u_0(\frac{\theta}{2})$ and $w(y) = v(2 \arctan y)((1 + y^2)/2)^{3/2}$ (see (6.3)). Since $u_0(a) = u_0(-a) = 0$ and $u'_0(-a) = u'_0(a) = 0$, we know that $w(\pm \tan a) = w'(\pm \tan a) = 0$ and $w(y) > 0$ for $y \in (-\tan a, \tan a)$. Hence

$$\begin{aligned} \int_{I_l} ((u_0)_{\theta\theta}^2 - 10(u_0)_\theta^2 + 9u_0^2) d\theta &= 16 \int_{-2a}^{2a} (v_{\theta\theta}^2 - \frac{5}{2}v_\theta^2 + \frac{9}{16}v^2) d\theta \\ &\geq \int_{-\tan a}^{\tan a} (w''(y))^2 dy > 0. \end{aligned}$$

□

The rest of the proof of the existence of a minimizer will be similar to the proof of the existence part of Theorem 1.

Using Lemma 1, we may assume that $u_i(0) = u_i(\pi)$ and $u_i(\pi/2) \geq u_i(3\pi/2)$. Let

$$\lambda_i = \sqrt{\frac{u_i(\frac{\pi}{2})}{u_i(0)}}, \text{ and } w_i(\theta) = \frac{\mathbf{T}_{\lambda_i} u_i}{\|\mathbf{T}_{\lambda_i} u_i\|_{L^\infty}}.$$

It is easy to see that $\{w_i\}$ is also a minimizing sequence and

$$w_i(0) = w_i(\frac{\pi}{2}) = \frac{\sqrt{u_i(0)u_i(\frac{\pi}{2})}}{\|\mathbf{T}_{\lambda_i} u_i\|_{L^\infty}}.$$

Furthermore $u_i(0) = u_i(\pi)$ implies $w_i(\pi) = w_i(0) = w_i(\pi/2)$. Since $\{w_i\}$ is a minimizing sequence with $\|w_i\|_{L^\infty} = 1$, Lemma 4 implies that $\|w_i\|_{H^2}$ is bounded. Hence $w_i \rightharpoonup w_0$ in H^2 . It follows from Sobolev embedding theorem that $w_i \rightarrow w_0$ in $C^{1,\alpha}$ for any $\alpha \in (0, \frac{1}{2})$. If $\min w_0 > 0$, then w_0 is a minimizer as being pointed out above. Therefore we only need to consider the case of $\min w_0 = 0$.

Suppose $w_0(\theta_0) = \min w_0 = 0$ for some $\theta_0 \in [0, \pi]$. Since $w_0(0) = w_0(\pi) = w_0(\pi/2) \geq w_0(3\pi/2)$, Lemma 5 implies that if $w_0(0) = 0$ then $w_0 \equiv 0$, which

contradicts with the fact that $\max w_0 = 1$. Hence $\theta_0 \neq 0$, $w_0(\theta_0) = w_0(\theta_0 + \pi) = 0$ and $w_0(\theta) > 0$ for $\theta \neq \theta_0, \theta_0 + \pi$.

For each i , suppose $w_i(\theta_i) = \min w_i(\theta)$. After rotating and up to a subsequence of w_i , we may assume that $\theta_0 = 0$, and $\theta_i = \theta_0 + \pi = \pi$, i.e. w_i attains its minimum at π , $w_i \rightarrow w_0$ and $w_0 > 0$ except at 0 and π .

Let

$$\tau_i = \sqrt{\frac{w_i(\frac{\pi}{2})}{w_i(0)}}, \quad \sigma_i(\theta) = \sigma_{\tau_i}(\theta), \quad \text{and} \quad v_i = \frac{\mathbf{T}_{\tau_i} w_i}{\|\mathbf{T}_{\tau_i} w_i\|_{L^\infty}}.$$

Again $\{v_i\}$ is a minimizing sequence and $\|v_i\|_{H^2} \leq C$ by Lemma 4. Therefore, up to a subsequence, $\{v_i\}$ converges weakly in H^2 to some $v_0 \in H^2$ and $v_i \rightarrow v_0$ in $C^{1,\alpha}$ for any $\alpha \in (0, \frac{1}{2})$.

It is easy to see that

$$v_i(0) = v_i(\frac{\pi}{2}), \quad \lim_{i \rightarrow \infty} \frac{v_i(\frac{3\pi}{2})}{v_i(\frac{\pi}{2})} = \frac{w_0(\frac{3\pi}{2})}{w_0(\frac{\pi}{2})} \leq 1, \quad \text{and} \quad v_i(\pi) \leq v_i(0).$$

Hence

$$(6.4) \quad v_0(0) = v_0(\frac{\pi}{2}), \quad \frac{v_0(\frac{\pi}{2})}{v_0(\frac{3\pi}{2})} \geq 1 \quad \text{and} \quad v_0(\pi) \leq v_0(0).$$

Applying Lemma 5 to v_0 , we see that $v_0(0) > 0$, otherwise $v_0(0) = v_0(\frac{\pi}{2}) = v_0(\pi) = v_0(\frac{3\pi}{2}) = 0$, contradiction. It follows from (6.4) that $v_0(\frac{\pi}{2}) \neq 0$, thus $v_0(\frac{3\pi}{2}) \neq 0$. Applying Lemma 5 again, we have $v_0(\pi) > 0$. For any $\theta \in [0, 2\pi] \setminus \{\frac{\pi}{2}, \frac{3\pi}{2}\}$, since w_i attains its minimum at π , we obtain

$$\begin{aligned} v_i(\theta) &= \frac{1}{\|\mathbf{T}_{\tau_i} w_i\|_{L^\infty}} w_i(\sigma_i(\theta)) (\tau_i^2 \cos^2 \theta + \tau_i^{-2} \sin^2 \theta)^{3/2} \\ &\geq \frac{1}{\|\mathbf{T}_{\tau_i} w_i\|_{L^\infty}} w_i(\pi) \tau_i |\cos \theta|^3 \\ &= v_i(\pi) |\cos \theta|^3 \rightarrow v_0(\pi) |\cos \theta|^3 > 0 \end{aligned}$$

Hence $v_0 > 0$, thus it is a minimizer.

To classify the extremal function $v(\theta)$, we again observe that

$$\begin{aligned} H_s^2(S^1) &= \{u \in H^2(S^1) : u > 0, \int_0^{2\pi} \cos 3\theta \cdot u^{-5/3} d\theta = \int_0^{2\pi} \sin 3\theta \cdot u^{-5/3} d\theta = 0 \\ &\quad \text{and} \quad \int_0^{2\pi} \cos \theta \cdot u^{-5/3} d\theta = \int_0^{2\pi} \sin \theta \cdot u^{-5/3} d\theta = 0\}. \end{aligned}$$

Thus $v(\theta)$ satisfies the Euler-Lagrange equation:

$$\begin{aligned} (6.5) \quad v_{\theta\theta\theta\theta} + 10v_{\theta\theta} + 9v &= \tau v^{-\frac{5}{3}} + \frac{a \cos 3\theta}{v^{\frac{8}{3}}} + \frac{b \sin 3\theta}{v^{\frac{8}{3}}} + \frac{c \cos \theta}{v^{\frac{8}{3}}} + \frac{d \sin \theta}{v^{\frac{8}{3}}} \\ &= \tau v^{-\frac{5}{3}} + \frac{A \cos(3\theta + \alpha_1)}{v^{\frac{8}{3}}} + \frac{B \cos(\theta + \alpha_2)}{v^{\frac{8}{3}}} \end{aligned}$$

where τ, a, b, c, d are Lagrange multipliers and $A = \sqrt{a^2 + b^2}$, $B = \sqrt{c^2 + d^2}$. We then follow the argument given in [4]. Multiplying both sides of (6.5) by $\cos(3\theta + \alpha_1)$

and $\cos(\theta + \alpha_2)$ respectively and integrating over $[0, 2\pi]$, we obtain that

$$(6.6) \quad \begin{cases} A \int_0^{2\pi} \frac{\cos^2(3\theta + \alpha_1)}{v^{8/3}} d\theta + B \int_0^{2\pi} \frac{\cos(3\theta + \alpha_1) \cos(\theta + \alpha_2)}{v^{8/3}} d\theta = 0 \\ A \int_0^{2\pi} \frac{\cos(3\theta + \alpha_1) \cos(\theta + \alpha_2)}{v^{8/3}} d\theta + B \int_0^{2\pi} \frac{\cos^2(\theta + \alpha_2)}{v^{8/3}} d\theta = 0. \end{cases}$$

It follows from the Hölder inequality that the determinant of the coefficient matrix of linear system (6.6) is always positive. Therefore (6.6) has only the trivial solution $A = B = 0$. Hence, v satisfies

$$v_{\theta\theta\theta\theta} + 10v_{\theta\theta} + 9v = \tau v^{-3},$$

which implies that the symmetric Q -curvature of $v^{-\frac{4}{3}}g_s$ is a constant. We hereby completes the proof of Theorem 3.

Remark 6. *It can be checked from the classification of constant 1-scalar curvature metrics that the symmetric Q -curvature of a constant 1-scalar curvature metric is a constant. Conversely, it is not clear whether it is true, even though we suspect that it is the case. Technically, we do have difficulty to classify the metrics with constant symmetric Q curvature. We conjecture that if u is a minimizer of $\inf_{H_s^2(S^1)} \mathbf{F}(u)$, then*

$$u(\theta) = c (\lambda^2 \cos^2(\theta - \alpha) + \lambda^{-2} \sin^2(\theta - \alpha))^{3/2},$$

for some $\lambda, c > 0$ and $\alpha \in [0, 2\pi)$, and the infimum is $\mathbf{F}(u) = 144\pi^4$.

Remark 7. *Direct computation shows that*

$$Q_g^\alpha = \frac{\alpha}{3} \Delta_g R_g^\alpha + (R_g^\alpha)^2,$$

which implies

$$\int Q_g^\alpha dS_g = \int (R_g^\alpha)^2 dS_g.$$

Therefore using Theorem 4 we obtain that with length fixed, the infimum of $\int (k_g)^2 dS_g$ is attained by metric with constant 4-scalar curvature. In addition, it can be shown that $\int Q_g dS_g$, therefore $\int (k_g)^2 dS_g$ is increasing and bounded above along the Q -curvature flow (1.12).

REFERENCES

- [1] Ai, J.; Chou, K.S.; Wei, J. Self-similar solutions for the anisotropic affine curve shortening problem. *Calc. Var. Partial Differential Equations* 13 (2001), no. 3, 311–337.
- [2] Chang, A. The Moser-Trudinger inequality and applications to some problems in conformal geometry, *Nonlinear PDE in Differential Geometry* (Park City, UT, 1992), 65–125, IAS/Park City Math. Ser., 2, Amer. Math. Soc., Providence, RI, 1996.
- [3] Chen, X. Calabi flow in Riemann surface revisited: A new point of view, *Intern. Math. Res. Notices*, No 6(2001), 275–297.
- [4] Chen, W.; Howard, R.; Lutwak, E.; Yang, D.; and Zhang, G. A generalized affine isoperimetric inequality, *J. Geom. Anal.* 14 (2004), no. 4, 597–612.
- [5] Fefferman, C.; Graham, R. Q -curvature and Poincaré metrics, *Math. Res. Lett.*, 9(2002), no 2-3, 139–151.
- [6] Hang, F. On the higher order conformal covariant operators on the sphere, preprint.
- [7] Li, J.; Zhu, M. Sharp local embedding inequalities, to appear in *Comm. Pure Appl. Math.*
- [8] Li, Y.Y. Remark on some conformally invariant integral equations: the method of moving spheres. *J. Eur. Math. Soc. (JEMS)* 6 (2004), no. 2, 153–180.
- [9] Lutwak, E. On some affine isoperimetric inequalities. *J. Differential Geom.* 23 (1986), no. 1, 1–13.

- [10] Ni, Y.; Zhu, M. One dimensional conformal metric flows,
<http://arxiv.org/abs/math.AP/0611319>
- [11] Sapiro, G.; Tannenbaum, A. On affine plane curve evolution, *J. Funct. Anal.* 119(1994), 79–120.
- [12] Schwetlick, H.; Struwe, M. Convergence of the Yamabe flow for "large" energies. *J. Reine Angew. Math.* 562 (2003), 59–100.
- [13] Wei, J. Personal communication.
- [14] Yang, P.; Zhu, M. On the Paneitz energy on standard three sphere. *ESAIM Control Optim. Calc. Var.* 10 (2004), no. 2, 211–223.
- [15] Ye, R. Global existence and convergence of Yamabe flow. *J. Differential Geom.* 39 (1994), no. 1, 35–50.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019,